



Regional Dynamic Model and Statistical Techniques: California Climate Certainties and Uncertainties

N.L. Miller^{1,2}, N.J. Schlegel^{2,1}, P.B. Duffy³, D.R. Cayan^{4,5},
H. Hidalgo⁴, J. Jin^{6,1}, H. Kanamaru⁴, M. Kanamitsu⁴, T. O'Brien⁷,
L.C. Sloan⁷, M.A. Snyder⁷, K. Yoshimura⁴

¹*Berkeley National Laboratory*

²*University of California, Berkeley*

³*Climate Central and Livermore National Laboratory*

⁴*University of California, San Diego*

⁵*U.S. Geological Service*

⁶*Utah State University, Logan*

⁷*San Diego University of California, Santa Cruz*

Sponsor: CEC – PIER Climate Change Scenarios Program Manager Guido Franco



Outline

- **Regional Downscaling Techniques for the California Assessments**
- **Intercomparing Downscaling Strengths and Weaknesses**
- **Quantification of Variables, Fluxes, and Time-Series**
 - **Temperature and Precipitation**
 - **Wind Magnitude and Direction**
 - **Pressure and Geopotential Height**
 - **Model Performance Summary**
- **Preparing the Climate Change Scenarios for the 2010 Assessment**
- **Next Steps and Concluding Remarks**



Dynamic and Statistical Downscaling Methods

- **Dynamic downscaling is a numerical weather prediction with complete meteorological equations resulting in most weather, and climate, variables and fluxes.**
- **Statistical downscaling is based on coarse-resolution predictors that lead to high-resolution predictands for *temperature and precipitation*.**
- **Dynamic downscaling requires a large amount of computational and data storage resources.**
- **Statistical downscaling is computationally inexpensive and many representations can be generated quickly.**



Approach

Downscaling Groups: Three Dynamic and One Statistical

- Berkeley Lab and UC-Berkeley: Miller, Schlegel, Jin
NCAR Weather Research and Forecasting Model with
(1) Rapid Update Cycle (**WRF-RUC**)
(2) Community and Model version 3 (**WRF-CLM3**)
- UC-Santa Cruz: Sloan, Snyder, O'Brien
ICTP Regional Climate Model Version 3 (**RegCM3**)
- UC-San Diego: Kanamitsu, Yoshimura, Kanamaru
NOAA Regional Spectral Model (**RSM**)
- UC-San Diego: Hidalgo, Dettinger, Cayan
Constructed Analogues Statistical Model (**CANA**)



Dynamic Model Land Surface Features

	Vegetation	Soil	Snow	Lake	River-Routing
WRF-CLM3	Up to 10 vegetation types in one grid Dynamic vegetation	10-layer soil Frozen soil	5-layer snow Liquid water within snow Variable snow density	10-layer lake Snow and ice on the lake included	A simple digital elevation model (DEM) to calculate water flow directions
WRF-RUC	One vegetation type in one grid cell NO Dynamic vegetation	6-layer soil Frozen soil	2-layer snow No liquid water within snow Fixed snow density	N/A	N/A
RegCM3-BATS	One vegetation type in one grid cell NO Dynamic vegetation	3-layer soil No Frozen soil	1-layer snow No liquid water within snow Fixed snow density	Hostetler lake model	N/A
RSM-Noah	One vegetation type in one grid cell NO Dynamic vegetation	4-layer soil	1-layer snow No liquid water within snow Fixed snow density	N/A	N/A



Constructed Analogues Statistical Technique (CANA)

Key assumptions:

- **Future climate patterns can be derived from linear combinations of the weather from a library of past observations.**
- **Coarse-resolution models are correct at fine-scale resolutions as well.**
- **Advantage - CANA downscaling techniques does not require specific climate forcing.**
- **Disadvantage - CANA assumes stationarity and will not capture significant changes in climate forcing.**



The Importance of Model Intercomparisons

- Intercomparisons provide quantitative evaluations of model and process performance compared to observations and other models.
- Intercomparisons are essential for understanding how model simulated projections of the future compare with the present.
- Intercomparisons allow for model advancements, leading to reduced errors, and improved model predictability.
- Improved model predictability will allow for *better decision making* of actions needed for climate change mitigation, adaptation, and coping strategies.




Model Standards for Intercomparing

- Each RCM was required to generate a 10 year historical simulation, 1 January 1980 to 31 December 1989.
- Each RCM used the same set of double nested domains and resolutions (A) western U.S. at 30-km and (B) CA at 10-km.
- Each RCM used the same set of external forcing, the NCAR/DOE Reanalysis II dataset for Initial and Lateral Boundary Conditions.
- Sea Surface Temperature (SST) updating was based on the AMIP Dataset, except for RSM which used the European Reanalysis 40 year dataset (ERA40).
- Each model saved a common set of specified variables, fluxes, mapped these onto common grids for analysis and followed the PCMDI protocols for IPCC AR4 Intercomparisons.



Model Domains and Resolutions




A - Western U.S. and Eastern Pacific Ocean, 30-km resolution, [139W21N x 104W51N]

B - California, Nevada, Eastern Pacific Ocean, 10-km resolution, [128W31N x 113W44N]




Summer Maximum Temperature

June - August






Difference relative to PRISM in Maximum 2-m air temperature during June-August.






Summer Minimum Temperature *June - August*






Difference relative to PRISM in Minimum 2-m Air Temperature during June-August.






Winter Maximum Temperature

December - February






Difference relative to PRISM in Maximum 2-m air temperature during December-February






Winter Minimum Temperature

December - February






Difference relative to PRISM in Minimum 2-m air temperature during December-February.






Cumulative Precipitation

November - March






Cumulative November – March Precipitation Correlation to PRISM.






Transect Analysis at 38.5 N and 34 N






Time-Space plots of Precipitation at 38.5N. West-East Section across the Russian and American River Basins






January 500hPa Geopotential Heights






500 hPa Geopotential Heights at 38.5N. West-East X-Section with Russian and American River Basins






500 hPa geopotential heights at 34N.




SEPTEMBER
1982


NARR





RSM



WRF CLM3




RegCM3




OCTOBER
1982


NARR





RSM



WRF CLM3




RegCM3




NOVEMBER
1982


NARR





RSM



WRF CLM3




RegCM3




DECEMBER
1982


NARR





RSM



WRF CLM3




RegCM3




**JANUARY
1983**


NARR





RSM



WRF CLM3




RegCM3




**FEBRUARY
1983**


NARR





RSM



WRF CLM3




RegCM3




**MARCH
1983**


NARR





RSM



WRF CLM3




RegCM3




MAY
1983


NARR





RSM



WRF CLM3




RegCM3




June
1983


NARR





RSM



WRF CLM3




RegCM3




JULY
1983


NARR





RSM



WRF CLM3




RegCM3




**AUGUST
1983**


NARR





RSM



WRF CLM3




RegCM3







Transect Analysis at 38.5 N and 34 N



Wind Direction Southern California July-Aug 1983 Latitude=34N



Wind Speed Southern California July-Aug 1983 Latitude=34N





Next Steps:

Regional Climate Change Projections

- **Simulation of the historical climate with the IPCC Global Climate Models as input forcing.**
- **Begin “time slice” simulations of early, mid, and late century California climates at 10 km resolution.**
- **Expand number of model ensemble members and further coordinate simulations with other groups.**
- **New studies have begin with a very high-resolution, multi-grid Coastal Ocean Circulation Model and WRF.**



Concluding Remarks

- **Downscaling is only as good as the large-scale forcing**
- **Downscaling has limitations, each model has unique and in numerous cases similar strengths and weaknesses.**
- **The Dynamic Models shown are state-of-the-art, yet still have problems simulating precipitation and other variables.**
- **The CANA technique performs at least as well as the dynamic models. BUT - very for a few variables: Temperature, Precipitation**
- **Dynamic Models do well simulating large-scale features such as the 500 mb geopotential heights, BUT subgrid parameterizations continue to be problematic.**